

[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[PopSet](#)[Taxonomy](#)[OMIM](#)[Books](#)Search for [Limits](#)[Preview/Index](#)[History](#)[Clipboard](#)[Details](#)[About Entrez](#)[Text Version](#)[Entrez PubMed](#)[Overview](#)[Help | FAQ](#)[Tutorial](#)[New/Noteworthy](#)[PubMed Services](#)[Journal Browser](#)[MeSH Browser](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[LinkOut](#)[Cubby](#)[Related Resources](#)[Order Documents](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)☐ 1: Org Lett 1999 Jul 15;1(1):117-20[Related Articles](#), [NEW Books](#), [LinkOut](#)

## Antihydrophobic cosolvent effects in organic displacement reactions.

Breslow R, Groves K, Mayer MU.

Department of Chemistry, Columbia University, New York, New York 10027, USA. rb33@columbia.edu

[formula: see text] Rates of reactions in water can be modified by the presence of antihydrophobic cosolvents such as ethanol and DMSO, which lower the energies of nonpolar surfaces. The rate effects reflect changes both in the solvation of nonpolar surfaces and also in the solvation of polar groups. The effects have been sorted out for some displacement reactions, revealing the geometry of an interesting branching reaction whose two paths show different antihydrophobic effects.

PMID: 10822546 [PubMed - indexed for MEDLINE]

[Privacy Policy](#)[Write to the Help Desk](#)[NCBI | NLM | NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act | Disclaimer](#)